

ORAL MELANOSIS: A CASE SERIES

Harvinder Kumar¹, Pankaj Chaturvedi²

ABSTRACT

There is very little information in literature about oral melanosis not associated with racial pigmentation or secondary to other syndromes. Various stimuli that can result in an increased production of melanin at the level of mucosa include trauma, hormones, radiation, and medications. Three such cases are reported in which stimulus for melanosis was mechanical trauma in form of smoking.

This case report may be cited as: Kumar H, Chaturvedi P. Oral Melanosis : A case series. J Postgrad Med Inst 2011; 25(4): 376-8.

INTRODUCTION

In our clinical practice we see many cases of oral pigmentary lesions the etiology of whom ranges from physiologic changes (e.g., racial pigmentation) to manifestations of systemic illnesses (e.g., Addison's Disease) and malignant neoplasms (e.g., melanoma and Kaposi's sarcoma). There is very little information in literature about oral melanosis not associated with racial pigmentation or secondary to other syndromes. We are reporting three such cases of oral melanosis which presented at different subsites in contrast to sites commonly seen in literature.

CASE REPORTS

First case was 38 years old female, who presented with brown pigmentation on ventral aspect of tongue and the history of tobacco use in form of chewing for a period of 6 years (Figure 1). Second case was 54 years old male, presented with brown pigmentation on dorsum of tongue and history of smoking for more than 15 years (Figure 2). Third case was 59 years old male, presented with brown black pigmentation on inner aspect of lip mucosa (Figure 3). Patient was chronic smoker for the last 20 years. All cases were clinically

asymptomatic. In all cases biopsy of the lesion was taken to confirm the diagnosis and to rule out oral melanoma as the lesions were present on subsites other than commonly seen in melanosis. Reassurance was given to all the cases. Lesion reduced after one year of stopping the smoking.

DISCUSSION

Oral melanosis is a benign focal pigmentary (brown or black) lesion of the oral cavity mucosa, though cases of melanosis have been reported on sinonasal, pharyngeal, conjunctival and laryngeal mucosa also. Oral pigmentation is caused by exogenous (foreign body implantation) and endogenous pigmentation. Primary pigments responsible for endogenous pigmentation include: melanin, melanoid, oxyhemoglobin, reduced haemoglobin, and carotene. Out of these pigments, melanin is the most common endogenous pigment which is a nonhemoglobin derived brown pigment and is produced by melanocytes present in the basal layer of the epithelium¹. Melanocytes were first identified in the oral epithelium by Becker in 1927 and a few years later they were isolated from samples of gingival tissue by Laidlaw and Cahn². During early intrauterine life, melanoblasts (precursor of melanocyte) migrate from the neural crest to the epidermis and hair follicles. Later they differentiate into dendritic cells and appear in head and neck region after approximately 10 weeks of gestation. Melanocytic dendrites reach a number of keratinocytes in the close vicinity, and through these dendrites, melanin is transported and transmitted to these epithelial cells, process called melanogenesis. The normal melanocytes of the oral mucosa have a small, round nucleus and a small amount of a clear cytoplasm. The number of melanocytes in the mucosa corresponds numerically to that of skin but with a reduced activity³. The color of the mucosa may be brown, blue, grey or black, depending on the amount and location of melanin in the tissues. Various stimuli

¹ Department of ENT, MAMC Agroha, Hisar, - India

² Department of ENT, Tata Memorial hospital, Mumbai - India

Address for Correspondence:

Dr. Harvinder Kumar
Associate Professor,
Department of ENT, MAMC Agroha, Hisar - India
E-mail: drharvinderent@gmail.com

Date Received: June 12, 2011

Date Revised: June 29, 2011

Date Accepted: July 18, 2011


Figure 1: A 38 year old female with Oral Melanosis



Figure 2: A 54 year old male with Oral Melanosis



Figure 3: A 59 year old male with Oral Melanosis



that can result in an increased production of melanin at the level of mucosa include trauma, hormones, radiation, and medications¹. Tobacco used as smoked and smokeless form induces oral mucosal changes in which intra-oral mucosal pigmentation is one of the clinical manifestations⁴. Smoker's melanosis occurs in up to 21.5% of smokers¹. Smoking may cause increased production of melanin, which may provide a biologic defence against the noxious agents present in tobacco smoke⁵. and it is hypothesized that melanin may play a role in detoxification of polycyclic amines nicotine and benzopyrene. The intensity of the pigmentation is related to the duration and amount of smoking. The changes are symptomless, it is not premalignant, and it seems that the pigmentation is reversible upon smoking cessation. A pigmentation prevalence of about 30%, is seen on the anterior attached gingiva⁶. The reason can be attributed to

fact that the total number of melanophores in the attached gingival is approximately 16 times greater than in the free gingival⁷. Women are more commonly affected than men usually after the third decade of life, which suggests a possible synergistic effect between the female sex hormones and smoking⁷. Clinically, the lesion usually presents as multiple brown pigmented macules less than 1 cm in diameter, localized mainly at the attached labial anterior gingival and the interdental papillae of the mandible. The microscopic appearance of melanosis is essentially similar to that seen in physiologic pigmentation or a melanotic macule. Oral melanosis is benign condition but biopsy should be performed if there is surface elevation or increased pigment intensity or pigmentation is in an unexpected site⁸. In a conclusion of study conducted on primary malignant melanoma of oral cavity in Japan with

special reference to melanosis, it was suggested that more than a third of the malignant melanomas develop from oral melanosis. The incidence of melanosis associated with oral melanoma was observed also among Caucasians as per reviewed by Chaudhary et al⁹ and in 50% of 12 melanomas collected by Moore and Martin, the pigmentation had been present for 1 to 5 years before melanoma developed¹⁰. Oral melanosis should be differentiated from ephelis, which is a circumscribed brown macule over skin that has been exposed to sunlight and shows increased melanin pigmentation in the basal-cell layer without an increase in the number of melanocytes^{11,12}. Since melanotic macules of the oral mucosa do not exhibit a significant increase in the number of melanocytes, the term lentigo is not considered appropriate for this type of lesion¹³.

CONCLUSION

Oral melanosis is benign pigmentary lesion of oral mucosa. It is seen more often in smokers. Though it is clinically asymptomatic, but remains matter of concern to rule out underlying malignancy. Biopsy should be considered if there is surface elevation or increased pigment intensity or pigmentation is in an unexpected site.

REFERENCES

1. Kauzman A, Pavone M, Blanas N, Bradley G. Pigmented lesions of the oral cavity: review, differential diagnosis, and case presentations. *J Can Dent Assoc* 2004;70:682-3.
2. Laidlaw GF, Cahn LR. Melanoblasts in the gum. *J Dent Res*1932;12:534-7.
3. Bologna JL, Orlow SJ. Melanocyte biology. In: Bologna JL, Jorizzo JL, Rapini RP, editors. *Dermatology*. 2nd ed. London: Mosby; 2003. p. 44.
4. Sayed M, Mirbod ,Stephen I, Ahing: Tobacco-associated lesions of the oral cavity: part I. Nonmalignant lesions. *J Can Dent Assoc* 2000;66:252-6.
5. Hedin C, Pindborg JJ, Daftary DK, Mehta FS. Melanin depigmentation of the palatal mucosa in reverse smokers: a preliminary study. *J Oral Pathol Med* 1992;21:440-4.
6. Axell T, Hedin CA. Epidemiologic study of excessive oral melanin pigmentation with special reference to the influence of tobacco habits. *Scand J Dent Res* 1982;90:434-42.
7. Patsakas A, Demetriou N, Angelopoulos A. Melanin pigmentation and inflammation in human gingiva. *J Periodontol* 1981;52:701-4.
8. Neville BW, Damm DD, Allen CM, Bouquet JE. *Oral and maxillofacial pathology*. 2nd ed. Toronto (ON): WB Saunders Company; 2002.
9. Chaudhry AP, Hampel A, Gorlin RJ. Primary malignant melanoma of oral cavity: a review of 105 cases. *Cancer* 1958;11:923-8.
10. Moore ES, Martin H. Melanoma of the upper respiratory tract and oral cavity. *Cancer* 1955;6:1167-76.
11. Pais S, Hegde SK , Bhat SS. Oral melanotic macule: a case report. *J Indian Soc Pedo Prev Dent* 2004;22:73-5.
12. Trodohl JN, Spragul WG. Benign and malignant mucosa: an analysis of 135 cases. *Cancer* 1970; 25:812-23.
13. Shapiro L, Zegarelli DJ. The solitary labial lentigo: a clinicopathologic study of 20 cases. *Oral Surg* 1971;31:87-92.

**GRANT SUPPORT, FINANCIAL
DISCLOSURE AND CONFLICT
OF INTEREST**

None Declared